

Unrecognized Cycloaddition Reactions of N-Alkyl- α - β -Unsaturated Imines Occurring in Biosystems and Their Biological Roles

Pradipta A., Tsutsui A., Latypova L., Chulakova D., Smirnov I., Kurbangalieva A., Tanaka K.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. Despite being fundamentally ubiquitous and important species in organic and bioorganic chemistry, the reactivities of N-alkyl- α - β -unsaturated imines have not been thoroughly explored due to their instability. Here, we describe novel reactivities of N-alkyl- α , β -unsaturated imines, which derived from α , β -unsaturated aldehydes (e.g., unsubstituted or substituted acrolein), alkylamines (e.g., amino-alcohols or diamines), and formaldehyde to produce 1,5-diazacyclooctanes, hexahydropyrimidines, and 1,3,5-triazacyclooctanes via formal [4 + 4], [4 + 2], and [4 + 2 + 2] cycloadditions in a stereocontrolled manner. We then also synthetically demonstrated that reaction between acrolein and biogenic amines (e.g., polyamines, noradrenaline, sphingosine), which ubiquitously exist in biosystems, proceed smoothly to give the corresponding 1,5-diazacyclooctanes. Finally, we also examined biological functions of the cycloaddition products and revealed for the first time their roles in oxidative stress mechanism and inhibition of amyloid- β (A β) 1-40 fibrillization.

<http://dx.doi.org/10.1007/s12668-016-0236-7>

Keywords

1,3,5-Triazacyclooctane, 1,5-Diazacyclooctane, Hexahydropyrimidine, Imino cycloaddition, N-Alkyl- α , β -unsaturated imine